

Symmetries of the resistance of mesoscopic samples in the quantum Hall regime

E. Peled,¹ Y. Chen,² E. Diez,^{2,*} D. C. Tsui,² D. Shahar,¹ D. L. Sivco,³ and A. Y. Cho³

¹Department of Condensed Matter Physics, The Weizmann Institute of Science, Rehovot 76100, Israel

²Department of Electrical Engineering, Princeton University, Princeton, New Jersey 08544, USA

³Bell Laboratories, Lucent Technologies, 600 Mountain Avenue, Murray Hill, New Jersey 07974, USA

(Received 24 December 2003; revised manuscript received 9 March 2004; published 22 June 2004)

The symmetry properties of the resistance of mesoscopic samples in the quantum Hall regime are investigated. In addition to the reciprocity relation, our samples obey new symmetries, that relate resistances measured with different contact configurations. Different kinds of symmetries are identified, depending on whether the magnetic field value is such that the system is above, or below, a quantum Hall transition. Related symmetries have recently been reported for macroscopic samples in the quantum Hall regime by Ponomarenko *et al.* [Solid State Commun. **130**, 705 (2004)], and Karmakar *et al.* (preprint cond-mat/0309694).

DOI: 10.1103/PhysRevB.69.241305

PACS number(s): 73.43.-f, 73.23.-b

Early studies of mesoscopic conductance fluctuations in the presence of a magnetic field (B) revealed an apparent puzzle:^{1,2} the pattern of fluctuations obtained from thin-film metallic samples exhibited no specific symmetry with respect to the reversal of B . These findings appeared puzzling because the conductivity of the samples was expected to follow the Onsager relations,^{3,4} $\sigma_{\alpha\beta}(B)=\sigma_{\beta\alpha}(-B)$, where α and β refer to coordinates, and therefore to have a clear symmetry upon B reversal. This apparent contradiction was soon settled by Benoit *et al.*,⁵ and Büttiker,^{6,7} who derived a general formula for the experimentally measured four-terminal resistance configuration and demonstrated that this resistance (or conductance) need not be symmetric with respect to the reversal of B . Instead, it should obey the reciprocity relation, stating that it will be symmetric with respect to the reversal of B and the simultaneous exchange of the current and voltage contacts:

$$R_{ij,kl}(B) = R_{kl,ij}(-B). \quad (1)$$

Here we use the standard notation of $R_{ij,kl}$ for V_{kl}/I_{ij} , where V_{kl} is the voltage difference between contacts k and l and I_{ij} is the current between contacts i and j .

These experiments, performed using metallic wires and loops, were limited to the low- B regime where Landau levels are unresolved. We present, in this Communication, an experimental study of the resistance of mesoscopic samples, designed to test their symmetries in the quantum Hall (QH) regime. We show that, in addition to the reciprocity relation, resistances measured near transitions between QH states exhibit symmetries that are not predicted by Büttiker's four-terminal resistance formula. These symmetries describe relations between resistances that are obtained with different contact configurations.

Our samples were prepared from two InGaAs/InAlAs wafers that contain a 200 Å quantum well. A two-dimensional electronic system is formed in the quantum well after illumination with a LED. Due to the short-range alloy scattering in our material the electronic system has a low mobility, limiting our study to the integer QH effect. The data were obtained from two samples, T2C and T1B. Sample

T2C was cooled twice (T2Cm2 and T2Cn2), and had a density and mobility of $n_s=1.15 \times 10^{11} \text{ cm}^{-2}$, $\mu=14\,000 \text{ cm}^2/\text{Vs}$ for both cooldowns. Sample T1B was cooled once (T1Bc2), and had $n_s=3.65 \times 10^{11} \text{ cm}^{-2}$, $\mu=44\,000 \text{ cm}^2/\text{Vs}$. The samples were wet etched to a Hall-bar geometry shown in Fig. 1(a). Special care was taken in the alignment of the metallic contacts to the Hall bars, to




FIG. 1. (Color online) (a) Geometry and contact numbering of our samples. The black areas in the figure represent Au–Ge–Ni alloyed contacts that were designed to reach the edges of the Hall bar. The Hall bar has a lithographic width of 2 μm, with a center-to-center distance of 4 μm between the longitudinal voltage contacts (2–3 and 6–5) and a distance of 24 μm between the current contacts (1–4). The four voltage contact-pairs that are used in the measurements are denoted by the corresponding resistances, R_L^t , R_L^b , R_H^l , and R_H^r ; (b) the correlation between the fluctuations of $R_{14,23}(B)$ and $R_{23,14}(-B)$; (c) the longitudinal resistance, $R_{14,23}(B)$ [R_L^t of Fig. 1(a)], of sample T2Cm2 at the vicinity of the $v=2-1$ transition, together with its reciprocity-equivalent resistance, $R_{23,14}(-B)$ [not shown in Fig. 1(a)].

ensure that voltage probes on opposite sides of a Hall bar will probe the same region of the sample. Due to the small size of our samples, their resistances display reproducible fluctuations whose magnitude and B -correlations near $B=0$ were used to extract the phase-coherence length, L_ϕ .⁸ For our samples $L_\phi = 1.1-1.3 \mu\text{m}$ at a temperature (T) of 10 mK, the T at which all of the data presented here were taken. Four-terminal resistance measurements were done using standard ac lock-in techniques with frequencies of 3–4 Hz and a current $I=1 \text{ nA}$, safely below $I=10 \text{ nA}$ where the fluctuations begin to diminish in size. B -field sweep rates were from 0.02 to 0.05 T/min, keeping the fluctuations independent of sweep rate.

We begin the presentation of our data with an experimental test of the reciprocity relation for our samples in the QH regime. In Fig. 1(c) we compare two reciprocity-equivalent resistances of sample T2Cm2 in the vicinity of the transition from the $\nu=2$ to the $\nu=1$ QH state ($\nu=2-1$ transition, where ν is the Landau level filling factor). Referring to the contact numbering in Fig. 1(a), the resistances compared in Fig. 1(c) are a longitudinal resistance, $R_{14,23}$, measured at positive B , and the resistance obtained after exchanging the current and voltage contacts, $R_{23,14}$, and taken at negative B polarity. Although the resistances are dominated by reproducible fluctuations we find that, in accordance with the reciprocity relation of Eq. (1), they have nearly the same pattern differing by only a small fraction of their amplitude. In order to quantify their similarity, we calculate the correlation between the fluctuations of the two resistances, normalized by the auto-correlation of each fluctuation pattern,^{9,10} and averaged over a B range of 0.2 T. In Fig. 1(b) we plot the results of the correlation calculation for the data of Fig. 1(c). The high values of the correlation, ranging from 0.63 to 0.88, attest to the similarity between $R_{14,23}(B)$ and $R_{23,14}(-B)$. Other resistances that are related via Eq. (1), such as $R_{14,62}(B)$ and $R_{62,14}(-B)$, were also found to have similarly high correlation values. While the reciprocity relation has been demonstrated before for the QH regime,¹¹ we are extending it here to samples whose resistances are dominated by mesoscopic fluctuations.

The main purpose of this Communication is to describe new symmetry properties, of four-terminal measurements, that are particular to the QH regime. These symmetries relate resistance measurements done with different contact configurations using samples in the mesoscopic regime. For simplicity we compare the two Hall and two longitudinal measurement configurations illustrated in Fig. 1(a). The current flows between contacts 1 and 4, and voltages are measured using the contact pairs 2–3 for the “top” longitudinal resistance (R_L^t) and 6–5 for the “bottom” longitudinal resistance (R_L^b). Similarly, we use the contact pairs 6–2 for the “left” and 5–3 for the “right” Hall resistances, R_H^l and R_H^r , respectively. For an ideal, macroscopic and homogeneous, sample there should be no difference between the “top” and “bottom”, or “left” and “right”, measurements and one would expect to find $R_L^t=R_L^b$ and $R_H^l=R_H^r$. In experiments this is rarely the case and, in general, each contact configuration yields a different result. In the following we show that, although our measurements can yield very different results depending on




FIG. 2. (Color online) (a) The correlation between the fluctuations of R_L^t and R_L^b ; (b) R_L^t and R_L^b vs B obtained from sample T2Cn2 in the vicinity of the $\nu=2-1$ transition. Inset: the same resistances near $B=0$.

the specific configuration, they are still linked by clear symmetry relations. These relations can involve the simultaneous exchange of contacts and B polarity, in a manner which is akin to that prescribed by the reciprocity relation.

Let us begin by considering the longitudinal configurations. In Fig. 2(b) we compare the R_L^t and R_L^b obtained from sample T2Cn2, near the $\nu=2-1$ QH transition. The style of the curves (solid or dashed line) corresponds to the voltage contacts used, as indicated in Fig. 1(a). The first observation we make from these data is that they can be divided into two B ranges according to the similarity between R_L^t and R_L^b . For $B < 3.016$ T, on the left side of the dashed line in the figure, R_L^t and R_L^b are virtually indistinguishable and their correlation, shown in Fig. 2(a), is close to unity. This is similar to the behavior we observe outside the QH regime, for the resistance fluctuations near $B=0$, see the inset of Fig. 2(b).

The picture changes dramatically as B is increased beyond 3.016 T. The R_L traces gradually begin to deviate from each other and eventually become uncorrelated. This can be clearly seen in Fig. 2(a), where we plot the correlation function between the two R_L 's. On the low- B side of the transition, for $B < 3.016$ T, the correlation is between 0.89 to 0.96, while on the high- B side, approximately above 3.3 T, it randomly fluctuates between 0.03 and 0.3, indicating the uncorrelated nature of the R_L 's. At the intermediate B range, 3.016–3.3 T, the correlation interpolates between these two regions.

The surprising result of our work is that, despite the uncorrelated appearance of the two R_L traces at the high- B side of the transition, they do not represent independent measurements. Instead we find that, upon the reversal of B , the “top” and “bottom” measurements are mapped onto each other. This is shown in Fig. 3(b) where we plot $R_L^t(B)$ together with $R_L^b(-B)$. The similarity of the traces is clearly improved and the correlation [Fig. 3(a)] is close to unity for the entire range of B . We thus identified a new symmetry for mesoscopic samples in the QH regime:

$$R_L^t(B) = R_L^b(-B). \quad (2)$$

This symmetry holds also for the low- B side of the transition, indicating that R_L in this region is symmetric in B .




FIG. 3. (Color online) (a) The correlation between the fluctuations of $R_L^t(B)$ and $R_L^b(-B)$; (b) $R_L^t(B)$ and $R_L^b(-B)$ of sample T2Cn2 in the vicinity of the $\nu=2-1$ transition.

To check whether this behavior is common to other QH transitions we repeated our measurements with a higher density sample, T1Bc2, allowing us to observe well-separated $\nu=4-3$, $3-2$, and $2-1$ QH transitions. The bottom pair of traces in each graph in Fig. 4 are measurements of R_L^t and R_L^b near each transition. The division into high- and low- B ranges is evident for all transitions studied. We have also verified the validity of Eq. (2) for these transitions.




FIG. 4. (Color online) R_L (bottom traces in each graph) and R_H (top traces) vs B obtained from sample T1Bc2 in the vicinity of the $\nu=4-3$ (a), $3-2$ (b), and $2-1$ (c) QH transitions.

The symmetry of Eq. (2) has been observed before in large, macroscopic, Hall-bar shaped samples, and has been attributed to the existence of a longitudinal density-gradient in the samples.¹² Recently, Ponomarenko *et al.* and Karmakar *et al.*¹³ reported on the observation of this symmetry and suggested a model that details how the symmetry originates from a density gradient. Due to a longitudinal density-gradient in their samples the “left” and “right” Hall resistances of the samples are not equal, $R_H^l \neq R_H^r$, and are instead found to be B -shifted with respect to each other. This then leads to the existence of a difference between the longitudinal resistances, R_L^t and R_L^b , since according to Kirchhoff’s law the Hall voltage difference is equivalent to a longitudinal voltage difference, or: $R_H^r - R_H^l = R_L^t - R_L^b$. In their model, the authors of Ref. 13 calculate the various R_L ’s and R_H ’s of the sample, with their explicit dependence on the density gradient, and show that for a linear density gradient the R_L ’s follow the symmetry of Eq. (2).

In the following paragraph we discuss the properties of R_H in our samples. We show that in our samples the differences between R_H^l and R_H^r do not amount to only a B -shift between the two measured traces. Each one of the R_H measurements displays a distinctly different pattern of fluctuations, that are related to the R_L fluctuations. The appearance of such fluctuation-dominated resistances that, nonetheless, obey the symmetry of Eq. (2) cannot be accounted for by a density gradient, indicating that our observations are not within the scope of the model suggested in Ref. 13.

In Fig. 4 we present, along with the R_L traces, the R_H measurements corresponding to each transition. First, we note that whenever $R_L^l = R_L^b$, as is the case for the low- B side of the data in the figure, $R_H^l = R_H^r$ must follow. Inspecting the R_H ’s in the figure we see that not only they are equal but they are also in their quantized state, $R_H^l = R_H^r = h/e^2$, where i is 2, 3, or 4. As B is increased through the transition, and the R_L ’s begin to separate, the R_H ’s, maintaining Kirchhoff’s law, separate as well and cease to be quantized. At this higher- B range, described in detail in a previous publication,¹⁴ we have found another kind of correlation, between the fluctuations of R_L and those of R_H :

$$R_L^t + R_H^l = R_L^b + R_H^r = h/(i-1)e^2. \quad (3)$$

We emphasize that these correlations appear only between specific R_L - R_H pairs, depending on the polarity of B : At positive B the correlated pairs are R_L^t - R_H^l , and R_L^b - R_H^r (the styles of the traces in Fig. 4 were chosen to highlight these correlations), while at negative B they are R_L^t - R_H^r and R_L^b - R_H^l . The switching of pairs at negative B is a result of our samples having R_H ’s that are antisymmetric with respect to the reversal of B , together with the B symmetry of R_L , Eq. (2).

Our findings can be summarized as follows: for a transition from a $\nu=i$ to a $\nu=i-1$ QH state the low- B side of the transition has $R_L^t = R_L^b$ and $R_H^l = R_H^r$. R_L^t and R_L^b are nonzero and exhibit fluctuations, while R_H^l and R_H^r are quantized to the value of the preceding QH plateau, h/e^2 . On the high- B side of the transition $R_L^t \neq R_L^b$ and $R_H^l \neq R_H^r$. $R_L^{t(b)}$ and $R_H^{l(r)}$ are anti-correlated, exhibiting fluctuations of equal magnitude and

opposite sign. Their sum equals the resistance value of the next QH plateau, $h/(i-1)e^2$. The B symmetry of the resistances can be neatly summarized by the following observation: For both R_L and R_H the effect of reversing the direction of B is equivalent to an exchange of the “top” and “bottom” voltage probes.

In our samples the QH series is terminated with a transition to an insulating phase as B is increased beyond the $\nu=1$ QH state. In the vicinity of this last transition, on which we have reported previously,¹⁵ the behavior of R_L and R_H is similar to that observed for the low- B side of the higher-LL transitions. Both R_L 's are dominated by reproducible fluctuations that are nearly equal, $R_L^t=R_L^b$, and the Hall resistances are quantized to h/e^2 , their value at the QH state preceding the transition, $R_H^l=R_H^r=h/e^2$.

The accepted theoretical model for describing transport in mesoscopic samples at the QH regime is based on Büttiker's four-terminal resistance formula extended to include the existence of electronic edge-states.¹⁶ The presence of edge states, whose chirality is determined by the polarity of B , may point in the direction of the origin of the symmetries presented in this Communication. However, our observed symmetries in the QH regime do not emerge from a straightforward application of the Büttiker multiprobe formula.

In the transport models of Streda, Kucera, and MacDonald¹⁷ and of Jain and Kivelson¹⁸ resistance fluctuations appear as a result of electrons scattering between the “top” and “bottom” edge states. When only one edge state is present, corresponding to conduction via the lowest Landau level alone, these models predict that the fluctuations will be limited to R_L , leaving R_H quantized. This situation is in agreement with our observations at the transition to the in-

sulating phase, but does not account for the quantized R_H on the low- B side of the higher Landau level transitions of $\nu=4-3$, 3-2, and 2-1.

In a recent numerical simulation¹⁹ Zhou and Berciu make use of Büttiker's formulation to describe the resistance in the QH regime as a result of an interplay between chiral edge currents and the tunneling between the “top” and “bottom” edges of the Hall bar. Their simulations reproduce many of the central features of our results, identifying a low- and high- B regions for all QH transitions, and predicting the symmetry of Eq. (2). According to their model, on the low- B side of the transitions transport is dominated by the presence of edge states, together with tunneling between the “top” and “bottom” sides of the sample, while on the high- B side transport is enabled only via the latter process, with no edge states or tunneling between the “left” and “right” sides of the sample.

To conclude, we presented an experimental study of the symmetries of the resistance of mesoscopic samples in the QH regime. We demonstrated new symmetries, relating longitudinal and Hall resistances of different contact configurations and B polarities. The resistances in the vicinity of all QH transitions were found to follow one of two possible sets of symmetries, one on the low- B and the other on the high- B side of the transitions.

The authors would like to thank M. Büttiker, Y. Oreg, A. Stern, and C. Zhou for useful discussions. This work is supported by the BSF and by the Koshland Fund. Y. C. is supported by the (U.S.) NSF. E. D. is supported by the Ramón y Cajal Program of the Spanish Minister of Science and Technology.

*Present address: Departamento de Física, Facultad de Ciencias, Universidad de Salamanca, E-37008 Salamanca, Spain.

¹C. P. Umbach, S. Washburn, R. B. Laibowitz, and R. A. Webb, Phys. Rev. B **30**, 4048 (1984).

²R. A. Webb, S. Washburn, C. P. Umbach, and R. B. Laibowitz, Phys. Rev. Lett. **54**, 2696 (1985).

³L. Onsager, Phys. Rev. **38**, 2265 (1931).

⁴H. B. G. Casimir, Rev. Mod. Phys. **17**, 343 (1945).

⁵A. D. Benoit, S. Washburn, C. P. Umbach, R. B. Laibowitz, and R. A. Webb, Phys. Rev. Lett. **57**, 1765 (1986).

⁶M. Büttiker, Phys. Rev. Lett. **57**, 1761 (1986).

⁷M. Büttiker, IBM J. Res. Dev. **32**, 317 (1988).

⁸P. A. Lee, A. D. Stone, and H. Fukuyama, Phys. Rev. B **35**, 1039 (1987).

⁹We define the normalized correlation between the fluctuations of any two resistance measurements, $R^{(1)}$ and $R^{(2)}$, as: correlation = $|\text{cor}(\delta R^{(1)}, \delta R^{(2)}) / \sqrt{\text{cor}(\delta R^{(1)}, \delta R^{(1)}) \times \text{cor}(\delta R^{(2)}, \delta R^{(2)})}|$. Here cor stands for the value of cross-correlation function at zero shift.

¹⁰W. J. Skocpol, P. M. Mankiewich, R. E. Howard, L. D. Jackel, D. M. Tennant, and A. D. Stone, Phys. Rev. Lett. **58**, 2347 (1987).

¹¹H. H. Sample, W. J. Bruno, S. B. Sample, and E. K. Sichel, J. Appl. Phys. **61**, 1079 (1987).

¹²K. v. Klitzing and G. Ebert, in *Two-Dimensional Systems, Heterostructures, and Superlattices*, Springer Series in Solid-State Sciences, edited by G. Bauer, F. Kuchar, and H. Heinrich, (Springer-Verlag, Berlin, 1984), Vol. 53, p. 242.

¹³L. A. Ponomarenko, D. T. N. de Lang, A. de Visser, V. A. Kulbachinskii, G. B. Galiev, H. Küntzel, and A. M. M. Pruisken, Solid State Commun. **130**, 705 (2004); B. Karmakar, M. R. Gokhale, A. P. Shah, B. M. Arora, D. T. N. de Lang, A. de Visser, L. A. Ponomarenko, and A. M. M. Pruisken, cond-mat/0309694 (unpublished).

¹⁴E. Peled, D. Shahar, Y. Chen, E. Diez, D. L. Sivco, and A. Y. Cho, Phys. Rev. Lett. **91**, 236802 (2003).

¹⁵E. Peled, D. Shahar, Y. Chen, D. L. Sivco, and A. Y. Cho, Phys. Rev. Lett. **90**, 246802 (2003).

¹⁶M. Büttiker, Phys. Rev. B **38**, 9375 (1988).

¹⁷P. Streda, J. Kucera, and A. H. MacDonald, Phys. Rev. Lett. **59**, 1973 (1987).

¹⁸J. K. Jain and S. A. Kivelson, Phys. Rev. Lett. **60**, 1542 (1988).

¹⁹C. Zhou and M. Berciu, cond-mat/0311365 (unpublished).